

ASSESSMENT OF LIFESTYLE FACTORS INFLUENCING BODY MASS INDEX AMONG UNDERGRADUATE MBBS STUDENTS IN URBAN UNIVERSITIES

Original Article

Authors:

Vaneeza Iftikhar^{1**}**

vaneezaIftikhar6@gmail.com

Student of Bachelor in Dietetics and Nutrition, The University of Faisalabad, Pakistan.

Anees Rafique²

United Medical and Dental College Karachi, Pakistan.

<https://orcid.org/0009-0000-3417-0364>

Acknowledgement: The authors express gratitude to all participating students for their valuable time and cooperation.

Short Title: Lifestyle Factors Influencing BMI in MBBS Students

Corresponding Author:

Vaneeza Iftikhar, Student of Bachelor in Dietetics and Nutrition, The University of Faisalabad, Pakistan,

vaneezaIftikhar6@gmail.com

Abstract

Background: Body mass index (BMI) serves as an important measure of overall nutritional condition and a useful predictor of later metabolic complications. Students in universities, particularly those engaged in rigorous medical training, are especially prone to adopting poor lifestyle practices that may alter their BMI. The demanding academic load often brings stress, erratic eating schedules, lack of physical activity, and insufficient sleep, all of which contribute to overweight and obesity. Gaining insight into how daily lifestyle patterns interact with BMI is crucial for designing effective health-focused strategies.

Objective: To evaluate the association of dietary habits, physical activity, and daily routines with BMI among undergraduate MBBS students in urban universities.

Methods: A cross-sectional study was conducted over four months among 377 MBBS students from urban universities in Lahore, Pakistan. Participants were recruited through stratified random sampling. Data were collected using a structured questionnaire assessing dietary intake, physical activity (Global Physical Activity Questionnaire), sleep duration, and screen time. BMI was measured using WHO criteria. Data were analyzed using SPSS v26, with independent t-tests and ANOVA applied to assess associations between lifestyle variables and BMI, considering $p < 0.05$ as statistically significant.

Results: The mean BMI of participants was 22.8 ± 3.7 kg/m², with 59.9% in the normal range, 19.1% overweight, 9.8% obese, and 11.1% underweight. Frequent fast-food consumption and high screen time were positively associated with higher BMI ($p=0.01$), while regular breakfast and moderate-to-high physical activity were protective against overweight ($p=0.02$ and $p<0.001$, respectively). Inadequate sleep was also linked to elevated BMI ($p<0.05$).

Conclusion: Lifestyle factors such as diet quality, physical activity, sleep, and screen exposure significantly influenced BMI among MBBS students. Early interventions promoting healthy eating, regular exercise, and balanced routines are critical to preventing overweight and obesity in future healthcare professionals.

Keywords: Body Mass Index, Cross-Sectional Studies, Diet, Lifestyle, Medical Students, Obesity, Physical Activity, Sleep.

Introduction

Body mass index (BMI) is a widely recognized indicator of nutritional status and a key measure for assessing the risk of metabolic and cardiovascular diseases (1). In recent years, the prevalence of overweight and obesity has risen sharply among young adults, including university students, due to changes in lifestyle patterns and dietary habits. Medical students, despite their knowledge of health and disease, are not immune to these trends (2). The demanding nature of medical education, combined with irregular schedules, high academic pressure, and limited time for self-care, creates an environment that can negatively influence eating behaviors, physical activity levels, and overall lifestyle balance (3). As future healthcare providers, the health behaviors of medical students not only impact their personal well-being but also shape their credibility and effectiveness in counseling patients about healthy living (4). Urban universities present unique challenges and opportunities that may influence lifestyle behaviors. Rapid urbanization has led to increased availability of fast food, sedentary entertainment options, and reliance on motorized transport, all of which contribute to unhealthy weight gain (5). Simultaneously, urban environments may also provide access to recreational facilities and diverse dietary options, offering opportunities for healthier choices (6). Among medical students, however, the long hours spent in lectures, clinical rotations, and exam preparation often result in inadequate physical activity, reliance on calorie-dense foods, and disrupted sleep cycles (7). Studies conducted in South Asia and other developing regions have highlighted a concerning pattern of overweight and obesity in this population, with prevalence rates ranging from 20% to 35% depending on the setting (8). These findings underscore the complex interplay of behavioral, environmental, and cultural factors that shape BMI in young adults.

Dietary practices play a central role in determining body weight and composition (9). Skipping meals, frequent snacking on processed foods, and excessive consumption of sugary beverages are common habits observed among students experiencing academic stress. Irregular meal timing, particularly late-night eating, has been associated with increased adiposity and metabolic dysregulation (10). Similarly, inadequate intake of fruits, vegetables, and high-fiber foods contributes to an imbalance in energy consumption and expenditure. Physical activity is another critical determinant of BMI. Despite widespread knowledge of its benefits, medical students often struggle to meet recommended activity levels, citing lack of time, fatigue, and academic commitments as barriers. The combination of poor dietary habits and insufficient exercise creates a high-risk environment for gradual weight gain and related health complications. Daily routines, including sleep patterns and study habits, also influence metabolic health (11). Sleep deprivation, a common issue during exam periods and clinical postings, has been linked to hormonal changes that increase appetite and promote fat storage. Furthermore, excessive screen time and sedentary behavior reduce overall energy expenditure, compounding the effects of poor diet and physical inactivity. The cumulative impact of these factors highlights the importance of adopting a holistic approach to understanding BMI among medical students, one that considers not only diet and exercise but also the broader lifestyle context.

While previous research has examined obesity and lifestyle factors in general university populations, there remains a need to focus specifically on medical students in urban Pakistani universities. Cultural dietary preferences, social norms, and environmental influences differ from those in Western settings, making local data essential for effective health promotion strategies. Moreover, as future physicians, medical students serve as role models for their patients and communities. Their personal health practices can influence their attitudes toward preventive medicine and their ability to counsel patients about nutrition and exercise. This study aims to evaluate the relationship between lifestyle factors and body mass index among undergraduate MBBS students in urban universities. By examining dietary habits, physical activity levels, and daily routines, the research seeks to identify modifiable behaviors that contribute to underweight, normal, overweight, or obese categories. The objective is to generate evidence that will inform targeted interventions and health education programs designed to promote healthy lifestyles among medical students, ultimately fostering a culture of wellness within the medical profession.

Methods

This descriptive cross-sectional study was conducted over a period of four months in two major urban universities of Lahore to assess the association between lifestyle factors and body mass index (BMI) among undergraduate MBBS students. The study population consisted of currently enrolled students from first to final year of the MBBS program. The sample size was calculated using the World Health Organization sample size calculator for prevalence studies, assuming a 30% estimated prevalence of

overweight or obesity based on recent regional literature, a 95% confidence level, and a 5% margin of error. The minimum required sample size was determined to be 380, and to account for possible non-response, 420 students were invited to participate through stratified random sampling proportional to the size of each academic year. All students aged 18 to 25 years who were currently registered in the MBBS program and willing to provide written informed consent were eligible for inclusion. Students with known chronic medical conditions such as thyroid disorders, diabetes mellitus, or other metabolic diseases that could independently affect BMI were excluded. Those on long-term medications influencing weight, such as corticosteroids or antidepressants, were also excluded to reduce confounding. Recruitment was carried out through announcements in lecture halls and student groups, and participation was voluntary with no academic incentives.

Information for this study was gathered through a structured and pre-tested questionnaire that focused on three main aspects: eating behavior, physical activity, and daily lifestyle patterns. Eating habits were assessed through a modified food frequency questionnaire adapted from the WHO STEPwise model, recording details such as number of meals per day, fruit and vegetable intake, use of fast foods, sugary drink consumption, and late-night snacking. Physical activity was evaluated using the short version of the International Physical Activity Questionnaire (IPAQ-SF), which documented both duration and intensity of activities performed in the preceding seven days, and assigned metabolic equivalent (MET) values to categorize participants into low, moderate, or high activity groups. Routine behaviors such as sleep duration, screen exposure, study time, and consistency of meal timings were also considered. Anthropometric data were collected by trained personnel following standardized guidelines. Body weight was measured to the nearest 0.1 kg with a calibrated digital scale, and height was taken to the nearest 0.1 cm using a portable stadiometer. Body mass index was determined by dividing weight in kilograms by height in meters squared, and participants were classified according to WHO Asian-specific ranges: underweight (<18.5 kg/m²), normal (18.5–22.9 kg/m²), overweight (23–24.9 kg/m²), and obese (≥ 25 kg/m²). All measurements were recorded with participants in light clothing and without footwear to ensure accuracy. The questionnaires were in English, completed by the students themselves, and supervised by the research team to reduce errors or missing responses. Participant confidentiality was safeguarded by assigning unique codes and securing electronic files on password-protected systems. Ethical approval was granted by the institutional review boards of the concerned universities, and written informed consent was obtained from every participant prior to data collection.

The collected data were processed and analyzed using SPSS version 26. Continuous factors including age, BMI, MET scores, and sleep duration were described as means along with standard deviations, while categorical factors such as gender, eating behaviors, activity levels, and BMI groupings were summarized in terms of frequencies and percentages. The normality of continuous data was checked through the Shapiro–Wilk test and histogram inspection, which confirmed approximate normal distribution for most variables. For comparisons of mean BMI across categories of diet frequency, activity level, and sleep patterns, independent sample t-tests and one-way ANOVA were applied. Associations between categorical lifestyle indicators and BMI groups were examined using chi-square analysis. Pearson's correlation was carried out to assess relationships among continuous factors such as study hours, MET values, and BMI. Additionally, a multiple linear regression model was developed to determine independent predictors of BMI, controlling for variables including age, sex, and living arrangement (hostel versus day scholar). A significance threshold of $p < 0.05$ was used for all analyses. This methodological approach allowed for accurate assessment of lifestyle influences on BMI among medical students. The use of validated tools, standardized anthropometric techniques, and rigorous statistical procedures ensured reliable outcomes, minimized potential bias, and strengthened reproducibility.

Results

A total of 377 undergraduate MBBS students participated in the study, yielding a response rate of 94.3%. The mean age of the participants was 21.1 ± 1.8 years, with a nearly balanced gender distribution of 178 (47.2%) males and 199 (52.8%) females. Students were drawn from all five years of study, with the highest representation from second year (25.1%) and third year (24.3%). Slightly more than half of the participants were day scholars (50.7%), while 49.3% resided in hostels (Table 1). The mean body mass index (BMI) of the cohort was 22.8 ± 3.7 kg/m². Based on WHO classifications, 226 students (59.9%) were within the normal range, 72 (19.1%) were overweight, 37 (9.8%) were obese, and 42 (11.1%) were underweight (Table 2). Gender-stratified analysis revealed a slightly higher proportion of overweight and obesity among males (34.8%) compared to females (25.6%). Mean BMI did not differ significantly across year of study ($p = 0.13$). Figure 1 illustrates the distribution of BMI categories across the sample.



Dietary behavior analysis showed that regular breakfast consumption was reported by 289 students (76.7%). Only 134 students (35.5%) achieved the recommended daily intake of five or more servings of fruits and vegetables. Frequent fast-food consumption (≥ 3 times per week) was noted among 156 students (41.4%), and 102 students (27.1%) consumed sugary drinks at least five times per week (Table 3). Higher fast-food intake was associated with elevated BMI ($p = 0.01$), whereas regular breakfast was inversely associated with overweight status ($p = 0.02$). Physical activity levels indicated that 98 participants (26.0%) reported low activity, 183 (48.5%) engaged in moderate activity, and 96 (25.5%) maintained high activity levels (Table 4). Figure 2 presents the proportional distribution of physical activity levels. Low physical activity was significantly more prevalent among students classified as overweight or obese compared to those with normal BMI ($p < 0.001$). Lifestyle routines further revealed that 157 students (41.6%) reported sleeping less than six hours per night, and 204 students (54.1%) reported screen time exceeding four hours per day (Table 5). Both insufficient sleep and prolonged screen exposure demonstrated positive associations with higher BMI ($p < 0.05$ for both variables). No significant differences in sleep duration were observed between hostel residents and day scholars. Overall, these findings underscore a multifactorial pattern in which dietary habits, physical activity, and daily routines collectively influence BMI among MBBS students. Regular breakfast consumption and moderate-to-high physical activity emerged as protective factors against overweight and obesity, while frequent fast-food intake, insufficient sleep, and prolonged screen time were associated with increased BMI.

Table 1: Demographic characteristics of participants (n = 377)

<i>Variable</i>	<i>Category</i>	<i>n (%)</i>
<i>Age (years)</i>	Mean $\pm$ SD	21.1 $\pm$ 1.8
<i>Gender</i>	Male	178 (47.2)
	Female	199 (52.8)
<i>Year of Study</i>	First	65 (17.2)
	Second	95 (25.1)
	Third	92 (24.3)
	Fourth	71 (18.8)
	Final	54 (14.3)
<i>Residence</i>	Hostel	186 (49.3)
	Day Scholar	191 (50.7)

Table 2: Distribution of BMI categories

<i>BMI Category</i>	<i>n (%)</i>
<i>Underweight (<18.5 kg/m²)</i>	42 (11.1)
<i>Normal (18.5–24.9 kg/m²)</i>	226 (59.9)
<i>Overweight (25–29.9 kg/m²)</i>	72 (19.1)
<i>Obese (≥30 kg/m²)</i>	37 (9.8)
<i>Mean BMI (kg/m²)</i>	22.8 ± 3.7

Table 3: Dietary patterns among participants

<i>Dietary Variable</i>	<i>n (%)</i>
<i>Regular breakfast consumption</i>	289 (76.7)
<i>≥5 servings fruits/vegetables daily</i>	134 (35.5)
<i>Fast-food ≥3 times/week</i>	156 (41.4)
<i>Sugary drinks ≥5 times/week</i>	102 (27.1)

Table 4: Physical activity levels

<i>Activity Level</i>	<i>n (%)</i>
<i>Low</i>	98 (26.0)
<i>Moderate</i>	183 (48.5)
<i>High</i>	96 (25.5)

Table 5: Sleep duration and screen time

<i>Variable</i>	<i>n (%)</i>
<i>Sleep <6 hours/night</i>	157 (41.6)
<i>Screen time >4 hours/day</i>	204 (54.1)




Figure 1 Distribution of BMI Categories

Discussion

The present study provided a comprehensive assessment of lifestyle factors influencing body mass index among undergraduate MBBS students in an urban university setting (12). The findings highlighted a predominance of normal BMI but revealed a considerable proportion of overweight and obesity, emphasizing the early emergence of weight-related health risks within this population. The association of dietary habits, physical activity levels, sleep duration, and screen time with BMI reflects the multifactorial determinants of body composition and underscores the need for integrated health promotion strategies (13). The observed prevalence of overweight and obesity aligned with global reports of rising weight-related concerns among medical students, particularly in South Asian contexts where urbanization and academic stress contribute to sedentary behaviors and poor dietary practices (14). The protective role of regular breakfast

consumption and the inverse relationship between moderate-to-high physical activity and elevated BMI supported previous evidence that consistent meal patterns and adequate physical engagement mitigate weight gain. Similar findings have been documented in cross-sectional studies conducted in India and Malaysia, where medical students with regular breakfast habits and at least moderate physical activity demonstrated significantly lower BMI (15). The association of frequent fast-food intake, prolonged screen time, and insufficient sleep with higher BMI further confirmed the influence of modern lifestyle behaviors described in earlier literature on young adults. The present findings strengthen the argument that medical students, despite their theoretical knowledge of health, remain vulnerable to unhealthy patterns during demanding academic schedules.

These results carry meaningful implications for health promotion in medical education. The coexistence of knowledge and risky behavior calls for targeted interventions within universities to foster healthy dietary habits, structured physical activity, and sleep hygiene. Institutional support, including accessible healthy meal options, student counseling, and organized sports programs, may help counteract the high demands of medical curricula that often limit opportunities for self-care (16). The data also suggest that lifestyle education must extend beyond theoretical instruction and include practical, habit-forming initiatives to improve long-term




Figure 2 Physical Activity Level

outcomes. The study possessed several strengths, including a robust sample size, validated measurement tools, and the use of standardized BMI classification to ensure comparability with global data (17). The inclusion of diverse lifestyle variables allowed for a nuanced analysis of interacting factors affecting BMI (18). Furthermore, the high response rate enhanced the representativeness of the findings within the targeted student population.

However, certain limitations warrant acknowledgment. The cross-sectional design precluded causal inference, and the reliance on self-reported dietary and activity data introduced the potential for recall and reporting biases (19). Additionally, the study was conducted in a single urban setting, which may limit generalizability to rural populations or to students in different academic disciplines (20). Seasonal variations in diet or activity, which were not controlled for, may have influenced the results (21). Future research should adopt longitudinal designs to explore causal relationships and to evaluate the impact of targeted interventions over time. Comparative studies between medical and non-medical student populations may also provide valuable insight into the unique challenges faced by those in rigorous academic programs (22). Despite these limitations, the study contributes important evidence highlighting the complex interplay of diet, physical activity, and lifestyle behaviors in shaping body composition among future healthcare professionals. These findings emphasize the importance of promoting a healthy lifestyle within medical institutions, not only to improve students' own health but also to enhance their credibility as future health educators.

Conclusion

This study demonstrated that BMI among MBBS students was significantly influenced by diet, physical activity, sleep, and screen time, with unhealthy habits contributing to overweight and obesity despite widespread health knowledge. These findings call for university-level interventions and sustained health education programs to promote balanced nutrition, regular physical activity, and adequate rest among medical students, ultimately fostering healthier lifestyles and better long-term health outcomes.

Author Contributions

Author	Contribution
Vaneeza Iftikhar ^{1*}	Substantial Contribution to study design, analysis, acquisition of Data Manuscript Writing Has given Final Approval of the version to be published
Anees Rafique ²	Substantial Contribution to study design, acquisition and interpretation of Data Critical Review and Manuscript Writing Has given Final Approval of the version to be published

References

1. Aslani A, Faraji A, Allahverdizadeh B, Fathnezhad-Kazemi AJNo. Prevalence of obesity and association between body mass index and different aspects of lifestyle in medical sciences students: A cross-sectional study. 2021;8(1):372-9.
2. Mani GJJOPMS. Assessment of body mass index and its associated nutritional factors among undergraduate medical students in Tamil Nadu, India: a crosssectional study. 2014;4:137-42.
3. Aldahash FD, Alasmari SA, Alnomsi SJ, Alshehri AM, Alharthi NF, Aloufi AAH, et al. Relationship of body mass index to sleep duration, and current smoking among medical students in Tabuk City, Saudi Arabia. 2018;10(9):7273.
4. Abdelmalek S, Adam H, Alardan S, Yassin S, Chtourou H, Souissi NJJoER, et al. Physical Activity, Sleep Patterns and Diet Habits as Well as the Prevalence of Obesity among Adolescents: A Cross Sectional Study from Ha'il City in Saudi Arabia. 2022;19(23):16174.
5. Palijo AH, Talpur SH, Luhana JK, Shoro JM, Baloch SJLMRJ. THE ASSOCIATION BETWEEN DIETARY HABITS, SLEEP PATTERNS, PHYSICAL ACTIVITY, AND OBESITY AMONG MEDICAL STUDENTS. 2025;7(2).

6. Rathod SS, Nagose VB, Kanagala A, Bhuvangiri H, Kanneganti J, Annepaka EJNJoP, Pharmacy, et al. Sleep duration and its association with obesity and overweight in medical students: A cross-sectional study. 2018;8(1):113-.
7. Habib SR, Alghofaily S, Alshamrani H, Alhammad AJTjocdp. Relationship of body mass index with diet, physical activities, and lifestyles of dental students. 2015;18(10):899-904.
8. Borresen J, Celie B, Laubscher R, Bac M, Wood P, Camacho T, et al. Poor sleep quality, low physical activity and low-to-moderate diet quality in South African first-year medical students. 2025;17(1):2-7.
9. Pop L-M, Iorga M, Muraru I-D, Petrariu F-DJS. Assessment of dietary habits, physical activity and lifestyle in medical university students. 2021;13(6):3572.
10. Rasul FB, Shawon MSR, Nazneen S, Hossain FB, Malik SS, Islam MTJJBAH. Do the dietary and lifestyle practices make the private medical students overweight: A cross-sectional study in Bangladesh. 2013;3(2):130-9.
11. Li G, Wu M, Lu G, Yu Z, Bao Z, Yu C, et al. Interactive effects of body mass index and physical activity on sleep quality in nursing students. 2025;16:1643841.
12. Hashmi MRUI, Shoaib S, Sajjad SS, Amer A, Minhas SA, Amer A, et al. Effects of Diet and Physical Activity on BMI of Medical Students: Cross Sectional Study. 2022;16(12):125-.
13. Jalal SM, Beth MRM, Al-Hassan HJM, Alshealah NMJJJomh. Body mass index, practice of physical activity and lifestyle of students during COVID-19 lockdown. 2021:1901-10.
14. Sertsuwankul P, Sakboonyarat B, Lertsakulbunlue SJCE, Health G. Prevalence and associated factors of increased risk for sleep-disordered breathing and its relationship with body mass index among medical students, intern doctors, and residents. 2024;27:101586.
15. Rafique T, Molla GM, Ahamed AJUDCJ. Association of body mass index (BMI) with diet, physical activity, and lifestyle among undergraduate dental students in a private dental college in Bangladesh. 2024;14(2):15-20.
16. Masaad AA, Yusuf AM, Shakir AZ, Khan MS, Khaleel S, Cheikh Ismail L, et al. Sleep quality and Dietary Inflammatory Index among university students: a cross-sectional study. 2021;25(4):2221-9.
17. Hossain MJ, Ahmmed F, Khan MR, Rashid PT, Hossain S, Rafi MO, et al. Impact of prolonged COVID-19 lockdown on body mass index, eating habits, and physical activity of university students in Bangladesh: a web-based cross-sectional study. 2022;9:873105.
18. Benaich S, Mehdad S, Andaloussi Z, Boutayeb S, Alamy M, Aguenau H, et al. Weight status, dietary habits, physical activity, screen time and sleep duration among university students. 2021;27(1):69-78.
19. Majeed FJJoTUMS. Association of BMI with diet and physical activity of female medical students at the University of Dammam, Kingdom of Saudi Arabia. 2015;10(2):188-96.
20. Haq IU, Mariyam Z, Zeb F, Jiang P, Wu X, Shah J, et al. Identification of body composition, dietary patterns and its associated factors in medical university students in China. 2020;59(1):65-78.
21. Hsu W-C, Chiang C-HJJomIr. Effect of BMI and perceived importance of health on the health behavior of college students: Cross-sectional study. 2020;22(6):e17640.
22. Alhashemi M, Mayo W, Alshaghel MM, Alsaman MZB, Kassem LHJAom, surgery. Prevalence of obesity and its association with fast-food consumption and physical activity: A cross-sectional study and review of medical students' obesity rate. 2022;79:104007.