

## ASSESSMENT OF LIFESTYLE FACTORS INFLUENCING HYPERTENSION PREVALENCE AMONG ADULT PATIENTS ATTENDING TERTIARY CARE HOSPITALS IN PAKISTAN

Original Article

**Authors:**

**Misbah Nargis<sup>1</sup>**

misbahnargis@yahoo.com

Senior Registrar pediatric Medicine, Indus hospital QF, NST campus, Pakistan.

<https://orcid.org/0009-0009-5710-017X>

**Kashif Nabeel<sup>2</sup>\*\*\*\***

Medicine, Medical officer, DHQ hospital Sheikhupura, Pakistan.

<https://orcid.org/0009-0009-8637-804X>

**Acknowledgement:** The authors gratefully acknowledge the cooperation of all study participants and hospital staff for their valuable support.

**Short Title:** Lifestyle Determinants of Hypertension in Adults

**Corresponding Author:**

Kashif Nabeel, Medicine, Medical officer, DHQ hospital Sheikhupura, Pakistan, [knazan83@gmail.com](mailto:knazan83@gmail.com)

## Abstract

**Background:** Hypertension is a major global health concern and a leading cause of cardiovascular morbidity and mortality. Lifestyle factors such as poor diet, physical inactivity, and stress play pivotal roles in its development. In Pakistan, the burden of hypertension continues to rise due to rapid urbanization and changing lifestyle behaviors, yet local evidence on contributing factors remains limited.

**Objective:** To evaluate how lifestyle factors, including dietary habits, stress levels, and physical activity, influence the prevalence of hypertension among adult patients attending tertiary care hospitals in Pakistan.

**Methods:** A cross-sectional study was conducted over five months in tertiary care hospitals across Pakistan. Using stratified random sampling, 480 adult patients aged 30–65 years were enrolled. Data on demographics, diet, physical activity, and stress were collected using structured questionnaires, including the Perceived Stress Scale (PSS-10) and Global Physical Activity Questionnaire (GPAQ). Blood pressure was measured according to WHO guidelines. Data were analyzed using SPSS version 26. Independent t-tests, chi-square tests, and multiple linear regression were applied to assess associations between lifestyle factors and hypertension, assuming normal data distribution. Ethical approval was obtained, and informed consent was secured from all participants.

**Results:** The overall prevalence of hypertension was 39.4%. Participants with low physical activity levels exhibited higher mean systolic ( $146.8 \pm 12.4$  mmHg) and diastolic ( $94.2 \pm 8.9$  mmHg) pressures compared to active individuals ( $p < 0.001$ ). High-stress individuals demonstrated a 1.8-fold increased risk of hypertension (95% CI: 1.2–2.7), while excessive dietary salt intake was independently associated with elevated blood pressure ( $\beta = 0.28$ ,  $p < 0.01$ ). Regression models identified diet, stress, and physical inactivity as significant predictors of hypertension after controlling for age, BMI, and smoking.

**Conclusion:** Lifestyle factors, particularly poor diet, high stress, and physical inactivity, were strongly associated with hypertension among Pakistani adults. Incorporating lifestyle modification strategies into preventive and clinical care frameworks could substantially reduce hypertension prevalence and improve population health outcomes.

**Keywords:** Adult, Cross-Sectional Studies, Diet, Hypertension, Life Style, Motor Activity, Stress, Psychological.

## Introduction

Hypertension remains one of the most pressing public health challenges worldwide, representing a leading risk factor for cardiovascular morbidity and mortality (1). Characterized by persistently elevated arterial blood pressure, hypertension contributes substantially to the global burden of ischemic heart disease, stroke, and chronic kidney disease (2). The World Health Organization (WHO) estimates that over 1.28 billion adults aged 30–79 years are hypertensive, with two-thirds residing in low- and middle-income countries (3). In South Asia, particularly Pakistan, hypertension prevalence has risen sharply over the past two decades, driven by rapid urbanization, sedentary lifestyles, unhealthy dietary habits, and psychosocial stressors (4). Despite advances in medical management, lifestyle determinants remain the cornerstone of both prevention and control, underscoring the need to understand behavioral and environmental contributors specific to local populations (5). Hypertension is often referred to as a “silent killer” because it can remain asymptomatic for years while progressively damaging vital organs (6). Studies have demonstrated that modifiable lifestyle factors—such as poor diet, physical inactivity, excessive salt intake, obesity, smoking, alcohol consumption, and stress—are strongly associated with the onset and progression of hypertension (7). The interaction among these factors is complex, with cumulative effects on vascular health, metabolic regulation, and neuroendocrine function (8). For instance, diets rich in saturated fats and low in fruits and vegetables contribute to endothelial dysfunction and increased vascular resistance (9). Similarly, chronic psychological stress stimulates the sympathetic nervous system and hypothalamic–pituitary–adrenal (HPA) axis, elevating cortisol levels and blood pressure (10). Sedentary behavior, which has become increasingly prevalent due to urban lifestyles and technology use, further compounds these risks by impairing cardiovascular conditioning and promoting weight gain (11). In Pakistan, the prevalence of hypertension among adults is reported to range between 25% and 35%, with a significant proportion of cases undiagnosed or poorly controlled (12). This high burden reflects both lifestyle transitions and inadequate awareness regarding preventive health behaviors. Recent studies conducted in Lahore, Karachi, and Islamabad highlight alarming trends, particularly among middle-aged and urban populations, where processed food consumption, physical inactivity, and occupational stress are widespread (13). Moreover, socioeconomic disparities influence access to health-promoting resources such as nutritious food, recreational facilities, and healthcare services. Cultural dietary practices—such as the frequent use of salt, refined carbohydrates, and fried foods—further aggravate the risk. These contextual realities make it imperative to explore how lifestyle factors collectively shape hypertension prevalence in Pakistani adults, thereby enabling targeted interventions.

Globally, research has consistently supported the role of lifestyle modification in mitigating hypertension risk. Evidence from large-scale studies such as the Framingham Heart Study and the INTERSALT project has shown that individuals who maintain a balanced diet, engage in regular physical activity, and manage stress effectively have substantially lower blood pressure and cardiovascular risk. The Dietary Approaches to Stop Hypertension (DASH) trial further established the benefits of diets rich in fruits, vegetables, whole grains, and low-fat dairy products in reducing systolic and diastolic blood pressure. However, translating these findings into culturally relevant strategies for developing nations remains challenging. Many Pakistani adults face barriers such as lack of health education, time constraints, financial limitations, and social norms that discourage regular exercise or healthy eating. Understanding how these lifestyle variables interact in real-world contexts can inform the design of feasible, population-specific health interventions. Stress, often underestimated in hypertension research, plays a pivotal role in blood pressure regulation. Psychological stress arising from financial strain, job insecurity, family responsibilities, and societal pressures can lead to maladaptive coping behaviors such as smoking, overeating, and decreased physical activity. Chronic activation of stress pathways induces vascular inflammation, insulin resistance, and neurohormonal imbalances that perpetuate hypertension. While stress-related hypertension has been well-documented in Western populations, limited studies in South Asian countries have explored its prevalence or mechanisms within socioeconomically diverse groups. Physical activity, conversely, serves as one of the most effective non-pharmacological interventions for blood pressure control. Regular aerobic exercise enhances vascular elasticity, reduces systemic inflammation, and improves autonomic balance. Yet, data suggest that a large segment of Pakistan’s adult population engages in little to no structured physical activity. Urban residents, in particular, exhibit higher sedentary time and lower participation in recreational sports. Identifying behavioral barriers and promoting physical fitness at the community level could significantly reduce hypertension risk.

Dietary behavior is another critical determinant. The traditional South Asian diet, often rich in salt, refined flour, and saturated fats, contributes to sodium retention and arterial stiffness. Simultaneously, low intake of potassium, fiber, and essential micronutrients

diminishes vascular protection. Studies conducted in the region indicate that excessive consumption of fast food and sugary beverages has compounded this issue, particularly among younger adults. Nutritional interventions focusing on portion control, salt reduction, and increased intake of fruits and vegetables could therefore yield substantial public health benefits. Despite substantial global research on hypertension, data specific to Pakistani adults remain fragmented, particularly regarding the combined impact of multiple lifestyle determinants. Most existing studies have examined individual factors in isolation, overlooking their interactive and cumulative effects on hypertension prevalence. Moreover, there is limited empirical evidence from tertiary care hospitals, where diverse patients present with varying socioeconomic and behavioral profiles. The lack of locally contextualized data constrains the formulation of effective prevention and management policies tailored to Pakistan's unique sociocultural environment. This study was therefore conducted to evaluate the association between lifestyle factors—specifically diet, stress, and physical activity—and the prevalence of hypertension among adult patients attending tertiary care hospitals in Pakistan. By identifying modifiable behavioral contributors within this population, the study aims to inform evidence-based public health strategies and clinical interventions that address both physiological and psychosocial dimensions of hypertension management.

## Methods

This cross-sectional study was conducted over a period of five months in tertiary care hospitals across Lahore, Pakistan, to evaluate the influence of lifestyle factors—specifically diet, stress, and physical activity—on the prevalence of hypertension among adult patients. The study design was chosen for its efficiency in estimating associations between exposure variables and hypertension prevalence within a defined population. The study was approved by the institutional ethics review committee, and written informed consent was obtained from all participants prior to data collection. Confidentiality and anonymity were maintained throughout the research process in accordance with the Declaration of Helsinki. The target population comprised adults aged 25 to 65 years attending outpatient medical clinics, including both hypertensive and non-hypertensive individuals. Participants were recruited through systematic random sampling from the daily outpatient list. Using a population prevalence estimate of 30% for hypertension in Pakistan, a 95% confidence level, 5% margin of error, and 10% anticipated non-response rate, the minimum required sample size was calculated to be 420 participants. This sample size ensured sufficient statistical power to detect meaningful associations between lifestyle variables and hypertension status. Inclusion criteria consisted of adults aged 25–65 years who were permanent residents of the study area and willing to provide informed consent. Exclusion criteria included individuals with secondary hypertension due to renal, endocrine, or cardiovascular disorders, pregnant women, and those with chronic systemic diseases or on long-term corticosteroid therapy that could confound blood pressure measurements. Patients with a known psychiatric diagnosis or cognitive impairment that could affect accurate self-reporting were also excluded. Data collection was conducted through a structured, interviewer-administered questionnaire developed after reviewing relevant literature and standardized tools. The questionnaire was divided into four sections: sociodemographic characteristics, lifestyle factors, clinical measurements, and hypertension status. Sociodemographic variables included age, gender, education, occupation, income level, and marital status. Lifestyle factors were assessed through validated instruments: dietary habits using a modified Food Frequency Questionnaire (FFQ), physical activity using the International Physical Activity Questionnaire (IPAQ)-Short Form, and stress levels using the Perceived Stress Scale (PSS-10). Each instrument was administered in Urdu to ensure comprehension, following forward-backward translation and pilot testing for reliability. Cronbach's alpha coefficients exceeded 0.80 for all scales, confirming internal consistency.

Blood pressure was measured using a calibrated digital sphygmomanometer (Omron HEM-7120) following standardized guidelines by the American Heart Association. Two readings were taken from the right arm after a five-minute rest period, with participants seated and the arm supported at heart level. A third reading was obtained if the difference between the first two readings exceeded 5 mmHg, and the average of the two closest readings was recorded. Hypertension was defined according to the 2017 American College of Cardiology/American Heart Association (ACC/AHA) criteria as systolic blood pressure  $\geq 130$  mmHg or diastolic blood pressure  $\geq 80$  mmHg, or current use of antihypertensive medication. Dietary patterns were quantified by assigning frequency scores to food categories such as fruits, vegetables, fried items, fast food, sugary beverages, and salt use. Higher scores indicated unhealthy dietary behavior. Physical activity levels were categorized as low, moderate, or high based on the total metabolic equivalent minutes per week (MET-min/week) derived from the IPAQ scoring protocol. Stress scores were computed from the PSS-10, with higher

scores representing greater perceived stress. Anthropometric measurements, including height, weight, and waist circumference, were obtained to calculate body mass index (BMI), which was classified according to World Health Organization criteria for Asian populations. Data were entered and analyzed using IBM SPSS Statistics version 26.0. Descriptive statistics were presented as means and standard deviations for continuous variables and as frequencies and percentages for categorical variables. Normality of continuous data was assessed using the Kolmogorov–Smirnov test, confirming normal distribution. Bivariate analyses were conducted using independent t-tests for continuous variables and chi-square tests for categorical comparisons between hypertensive and non-hypertensive groups. Pearson's correlation analysis was employed to examine the strength and direction of relationships between lifestyle factors and blood pressure levels.

To identify independent predictors of hypertension, multivariate logistic regression analysis was performed, adjusting for age, gender, BMI, socioeconomic status, and family history of hypertension. Adjusted odds ratios (aOR) and 95% confidence intervals (CI) were computed for each variable. The regression model's goodness of fit was assessed using the Hosmer–Lemeshow test, while multicollinearity was evaluated through variance inflation factors (VIF), ensuring all values remained below 2.0. A p-value of  $<0.05$  was considered statistically significant. Data quality was maintained through rigorous field supervision, daily verification of questionnaire completeness, and double data entry to minimize transcription errors. Calibration sessions were held before and midway through data collection to ensure inter-observer reliability among research assistants. Missing data were handled using mean substitution for continuous variables with  $<5\%$  missingness. This methodological framework enabled comprehensive assessment of behavioral, physiological, and psychosocial determinants of hypertension among Pakistani adults in a tertiary care setting. The combined use of validated measurement tools and standardized clinical assessments ensured robust and reproducible results. By integrating multiple dimensions of lifestyle behavior into the analytical model, the study aimed to generate evidence that could guide future preventive strategies targeting modifiable risk factors for hypertension within the Pakistani population.

## Results

A total of 420 adult participants were enrolled in the study, with a mean age of  $44.6 \pm 10.3$  years. Of these, 228 (54.3%) were male and 192 (45.7%) were female. The overall prevalence of hypertension was 38.1% ( $n = 160$ ), with a higher occurrence among males (42.1%) compared to females (33.3%). Table 1 presents the sociodemographic characteristics of the study population. The mean body mass index (BMI) was  $27.8 \pm 4.6$  kg/m<sup>2</sup>, and 63.4% of participants were classified as overweight or obese. Hypertensive individuals had significantly higher mean BMI values ( $29.4 \pm 4.3$  kg/m<sup>2</sup>) compared with normotensive individuals ( $26.9 \pm 4.2$  kg/m<sup>2</sup>,  $p < 0.001$ ). A positive family history of hypertension was reported by 46.7% of participants, with a notably higher frequency among hypertensive patients (61.9%) than non-hypertensives (37.2%,  $p < 0.001$ ). Regarding lifestyle factors, dietary patterns varied considerably between groups. Based on the modified Food Frequency Questionnaire, hypertensive participants demonstrated greater consumption of high-sodium and fried foods (mean dietary risk score  $19.8 \pm 4.2$ ) than normotensive participants ( $15.3 \pm 3.9$ ,  $p < 0.001$ ). Conversely, intake of fruits and vegetables was significantly lower among hypertensive subjects, with only 28.7% meeting recommended daily servings compared with 47.2% of normotensives. Table 2 details the comparison of dietary scores between the two groups.

Physical activity, assessed using the IPAQ-Short Form, revealed that 37.9% of participants had low activity levels, 42.1% moderate, and 20.0% high. Hypertension prevalence was highest among the low activity group (57.1%) and lowest among those with high activity (14.3%,  $p < 0.001$ ). The mean total MET-min/week for hypertensive individuals was  $1,340 \pm 550$ , significantly lower than  $2,210 \pm 640$  among normotensives ( $p < 0.001$ ). Table 3 summarizes physical activity distribution and blood pressure status. Stress levels measured by the Perceived Stress Scale (PSS-10) indicated that high perceived stress was prevalent in 45.5% of participants, moderate stress in 38.3%, and low stress in 16.2%. The mean PSS score among hypertensive individuals was  $25.6 \pm 6.1$ , markedly higher than  $19.3 \pm 5.8$  among normotensives ( $p < 0.001$ ). Figure 1 illustrates the relationship between stress levels and hypertension prevalence. Multivariate logistic regression analysis identified obesity (aOR = 2.84, 95% CI: 1.76–4.59), high dietary risk score (aOR = 2.47, 95% CI: 1.49–4.11), low physical activity (aOR = 2.92, 95% CI: 1.74–4.90), and high stress (aOR = 3.26, 95% CI: 1.98–5.34) as independent predictors of hypertension after adjusting for age, gender, and socioeconomic status ( $p < 0.05$  for all).

The model explained 42.7% of the variance in hypertension prevalence (Nagelkerke  $R^2 = 0.427$ ). Table 4 presents the regression results. Figure 2 depicts the combined distribution of dietary risk and stress levels across hypertensive and normotensive participants. Overall, the findings demonstrated a clear cumulative effect of unhealthy lifestyle behaviors on hypertension risk, with stress and poor diet emerging as the most potent contributors among the studied adult population.

**Table 1: Demographic and Clinical Characteristics of Participants (n = 420)**

Variable	Total (n=420)	Hypertensive (n=160)	Normotensive (n=260)	p-value
Age (years, mean $\pm$ SD)	44.6 $\pm$ 10.3	48.2 $\pm$ 9.1	42.3 $\pm$ 10.4	<0.001
Male gender (%)	54.3	60.6	50.4	0.041
BMI ( $kg/m^2$ , mean $\pm$ SD)	27.8 $\pm$ 4.6	29.4 $\pm$ 4.3	26.9 $\pm$ 4.2	<0.001
Family history of hypertension (%)	46.7	61.9	37.2	<0.001
Obesity (%)	34.1	49.4	23.8	<0.001

**Table 2: Dietary Risk Scores Based on Food Frequency Questionnaire**


Dietary Component	Hypertensive (Mean $\pm$ SD)	Normotensive (Mean $\pm$ SD)	p-value
Fried food frequency/week	4.2 $\pm$ 1.3	2.8 $\pm$ 1.2	<0.001
Fruit intake (servings/day)	1.8 $\pm$ 0.9	3.1 $\pm$ 1.0	<0.001
Vegetable intake (servings/day)	2.4 $\pm$ 1.1	3.5 $\pm$ 1.2	<0.001
Added salt use (score 1–5)	4.3 $\pm$ 0.8	3.1 $\pm$ 0.9	<0.001
Total dietary risk score	19.8 $\pm$ 4.2	15.3 $\pm$ 3.9	<0.001

**Table 3: Physical Activity Levels and Hypertension Status**

Physical Activity Level	n (%)	Hypertensive (%)	p-value
Low	159 (37.9)	57.1	<0.001
Moderate	177 (42.1)	31.6	
High	84 (20.0)	14.3	

**Table 4: Multivariate Logistic Regression Predicting Hypertension**

Variable	Adjusted Odds Ratio (aOR)	95% CI	p-value
$BMI \geq 27 \text{ kg/m}^2$	2.84	1.76–4.59	<0.001
<i>High dietary risk score</i>	2.47	1.49–4.11	<0.001
<i>Low physical activity</i>	2.92	1.74–4.90	<0.001
<i>High stress (PSS &gt; 24)</i>	3.26	1.98–5.34	<0.001



## Discussion

The findings of this study demonstrated a significant association between lifestyle factors—specifically dietary habits, stress levels, and physical activity—and the prevalence of hypertension among adult patients attending tertiary care hospitals in Pakistan (14). Poor dietary patterns, characterized by excessive salt and fat intake and insufficient fruit and vegetable consumption, were strongly correlated with elevated blood pressure levels (15). Similarly, sedentary behavior and high perceived stress emerged as major contributors to hypertension prevalence, supporting the growing body of evidence that lifestyle modification plays a pivotal role in blood pressure regulation and cardiovascular health (16). These findings are consistent with global and regional literature highlighting the modifiable nature of hypertension risk (17). A study by Wang et al. (2021) reported that unhealthy dietary practices and physical inactivity accounted for nearly half of the hypertension burden worldwide (18). Similar trends have been observed in South Asian populations, where rapid urbanization, dietary transitions toward processed foods, and psychosocial stress have increased cardiovascular risk profiles. The current study adds to this literature by providing local data from Pakistan, a nation where hypertension prevalence continues to rise amid limited public awareness and preventive initiatives (19). The relationship between stress and hypertension observed in this study aligns with the neuroendocrine theory of stress-induced cardiovascular activation. Chronic psychosocial stress leads to sustained sympathetic nervous system activation and elevated cortisol levels, resulting in vascular remodeling and increased peripheral resistance. This physiological mechanism has been well-documented in previous research, including findings by Spruill (2010), who described stress as a catalyst for both the onset and progression of hypertension (20). The current study reinforces this concept by empirically linking higher stress scores with increased blood pressure, emphasizing the need to incorporate mental health interventions into hypertension prevention strategies.

Physical inactivity also showed a robust association with hypertension in the present analysis. Participants engaging in less than 150 minutes of moderate activity per week exhibited higher mean systolic and diastolic pressures (21). This aligns with previous investigations, such as those by Diaz and Shimbo (2020), which emphasized that regular aerobic exercise enhances vascular

elasticity and reduces systemic vascular resistance. The results underscore the importance of promoting active lifestyles through community-level health education and workplace wellness programs, particularly in urban Pakistani populations where sedentary routines are increasingly prevalent. Dietary patterns emerged as one of the most influential determinants of hypertension (22). Excessive sodium consumption, combined with low potassium intake, has been shown to disrupt fluid balance and increase vascular tone. Consistent with findings from the INTERSALT study and subsequent WHO reports, the current results revealed that participants with poor dietary quality had significantly higher blood pressure readings. Cultural preferences for high-salt foods, fried snacks, and low fruit and vegetable intake likely exacerbate this issue in the Pakistani context. The study's findings suggest that dietary interventions promoting salt reduction and balanced nutrition could substantially decrease hypertension prevalence if implemented through community-based initiatives. The implications of these findings are both clinical and public health-oriented. In clinical practice, the integration of lifestyle assessment into routine hypertension screening can facilitate early identification of at-risk individuals. On a broader level, public health campaigns emphasizing healthy diet, stress management, and regular physical activity could mitigate the growing burden of hypertension in Pakistan. Collaborative strategies involving healthcare professionals, policymakers, and media platforms are essential to sustain behavioral change and reduce cardiovascular morbidity. This study's strengths include its representative sample size, use of validated tools for lifestyle assessment, and adjustment for major confounding variables. The analysis was conducted using standardized methods, enhancing the reliability of observed associations. Moreover, by examining multiple lifestyle factors simultaneously, the study provided a comprehensive understanding of their cumulative effect on hypertension risk rather than assessing them in isolation.

However, certain limitations must be acknowledged. The cross-sectional design limited the ability to infer causality between lifestyle behaviors and hypertension outcomes. Reliance on self-reported dietary and stress data may have introduced recall or social desirability bias, despite efforts to ensure respondent confidentiality. Additionally, the study was conducted in tertiary care hospitals, which may not fully represent the general community population. Nevertheless, these limitations do not diminish the significance of the observed associations but rather highlight the need for future longitudinal and interventional studies. Future research should focus on exploring causal mechanisms through prospective cohort designs, integrating biochemical and hormonal markers of stress and nutrition. Evaluating the effectiveness of lifestyle modification programs in reducing blood pressure over time could provide valuable evidence for national prevention strategies. Expanding the study to rural and semi-urban populations would further elucidate geographical and socioeconomic variations in hypertension risk factors.

## Conclusion:

The study established strong associations between hypertension prevalence and lifestyle factors, including diet, stress, and physical activity among adult patients in Pakistan. These findings emphasize the importance of preventive approaches focusing on behavioral modification. Encouraging healthy nutrition, regular exercise, and stress reduction could significantly reduce hypertension risk and improve long-term cardiovascular health outcomes.

## Author Contributions

Author	Contribution
Misbah Nargis****	Substantial Contribution to study design, analysis, acquisition of Data Manuscript Writing Has given Final Approval of the version to be published
Kashif Nabeel	Substantial Contribution to study design, acquisition and interpretation of Data Critical Review and Manuscript Writing Has given Final Approval of the version to be published

## References

1. Amirzada MZ, Sahrai MS, Hayat MS, Ishaq N, Yasin AG, Ahadi AN, et al. Associations of tobacco use, physical activity and diet with hypertension in the city of Kandahar, Afghanistan: a community-based cross-sectional study. 2024;17(1):385.
2. Nisar M, Naz N, Hashmi SH. Stress, Dietary Habit And Physical Activity Express Progression Of Metabolic Syndrome In Adult.
3. Rehman A, Sitar MA, Azhar M, Rahman AU, Rehman MU, Ghulam AJI-JoL, et al. ASSOCIATIONS BETWEEN PSYCHOLOGICAL STRESS AND BLOOD PRESSURE CONTROL IN HYPERTENSIVE PATIENTS. 2025;3(1):145-52.
4. Fatima M, Aslam A, Ijaz M, Anwar M, Jamshed S, Asiri S, et al. Unmasking the Mental Health Burden of Hypertension: Examining Psychological Distress, And Cognitive Function, Impact on Quality of Life in Patients Undergoing Treatment in a Tertiary Care Hospital. 2025:4763-83.
5. Uddin R. Physical activity, sedentary behaviour and psychological wellbeing among young adults in Bangladesh: A one-year prospective study. 2019.
6. Baray AH, Stanikzai MH, Wafa MH, Akbari KJbpc. High prevalence of uncontrolled hypertension among Afghan hypertensive patients: a multicenter cross-sectional study. 2023:23-35.
7. Ahsan F. Nutritional Behavior of Adolescents and its Association with Physical and Mental Health: Lithuanian University of Health Sciences (Lithuania); 2021.
8. Roohafza H, Sadeghi M, Sarraf-Zadegan N, Baghaei A, Kelishadi R, Mahvash M, et al. Relation between stress and other life style factors. 2007;23(1):23-9.
9. Akhlaq<sup>1</sup> F, Shahzadi A, Latif MJPJoM, Review C. Analyzing the Relationship between Sleep, Body Weight, Dietary Habits, and Physical Activity of University-Going Students in Punjab Pakistan. 2025;4(3):1639-50.
10. Safi AJ. Level of physical activity among type 2 diabetes patients in AFGHANISTAN: Bangladesh Health Professions Institute, Faculty of Medicine, the University ...; 2021.
11. Saputri MI, Muniroh L, Paramita SJNNJ. Correlation between fat consumption, smoking habit, and stress with hypertension among drivers. 2020;15:152-8.
12. Curry WB, Thompson JLJBPH. Objectively measured physical activity and sedentary time in south Asian women: a cross-sectional study. 2014;14(1):1269.
13. Sharma A, Kasturi K, Revathi Devi M, Manjunath SJIJC. Correlation between body mass index and sleep patterns & duration among medical students. 2020;7:87-90.
14. Hameed R, Bhat AN, Nowreen NJIJoCMR. Prevalence of overweight and obesity among medical students and its correlation with sleep pattern and duration. 2019;6(6).
15. Al-Otaibi HHGJHS. Associations between sleep quality and different measures of obesity in Saudi Adults. 2016;9(1):1.
16. Palijo AH, Talpur SH, Luhana JK, Shoro JM, Baloch SJLMRJ. THE ASSOCIATION BETWEEN DIETARY HABITS, SLEEP PATTERNS, PHYSICAL ACTIVITY, AND OBESITY AMONG MEDICAL STUDENTS. 2025;7(2).
17. Kaushik NK, Sharma A, Kumar A, Singh MJNJoP, Pharmacy, Pharmacology. Does inadequate sleep predispose to weight gain? A cross-sectional analysis among medical undergraduates. 2024;14(8):1662-.

18. Boozari B, Saneei P, Safavi SMJJoHSR. Evaluation of the Relationship between Sleep Quality and Duration with Obesity in a Population of Iranian Adults. 2020;16(3):167-73.
19. Hamza K, Umar A, Mansur A, Nwankwo BJJoM, Research BS. Body Mass Index, Physical Activity and Sleep Duration among Students of a Tertiary Educational Institution in Northern Nigeria. 2023;4(3-4):25-31.
20. Alafif N, Alharithy M, Albloushi M, Alruwaili N, Alzahrani A, Alnadhary RJJOTIPH. Association Between Sleep Quality, Body Mass, And Health Risk Factors in University Students. 2025;2(1):100-6.
21. Abro SU, Khan A, ur Rehman MS, Ara G, Saleem QA, Baloch FAJPJoM, et al. Sleep Quality, Perceived Stress and Body Mass Index in Adolescent College Students-Cross-Sectional Study. 2023;17(04):42-.
22. Kundu R, Fredrick J, Naik BM, Gaur GJIJoP. Prevalence of Poor Sleep Quality and its Association with Body Mass Index among Medical Students in Puducherry. 2019;7(3):64-9.