

ASSOCIATION BETWEEN ACADEMIC STRESS, SLEEP DISTURBANCE, AND DIETARY HABITS AMONG UNDERGRADUATE MBBS STUDENTS IN PAKISTAN

Original Article

Authors:

Jahanzaib Ali

www.ali24448@gmail.com

MBBS (3rd year) Student, Liaquat university of medical and Health sciences LUMHS Jamshoro, Pakistan.

<https://orcid.org/0009-0008-3282-7885>

Fatima

MBBS, Punjab medical college Faisalabad, Oral experts group Australia.

<https://orcid.org/0009-0006-0062-4931>

Majida Khan

Assistant Professor, LUMHS Jamshoro, Pakistan.

Acknowledgement: The authors gratefully acknowledge the cooperation of all participating medical students for their valuable contribution to this study.

Short Title: Academic Stress, Sleep, and Diet Among MBBS Students

Corresponding Author:

Jahanzaib Ali, MBBS (3rd year) Student, Liaquat university of medical and Health sciences LUMHS Jamshoro, Pakistan,

www.ali24448@gmail.com

Abstract

Background: Medical education imposes high academic demands, often resulting in elevated stress levels that adversely affect sleep quality and dietary habits. Among medical undergraduates, these lifestyle disruptions may contribute to both physical and psychological health concerns. Understanding these associations is essential for promoting well-being and academic performance among future healthcare professionals.

Objective: To assess the relationship between academic stress levels, sleep quality, and dietary behaviors among MBBS students in Pakistan, and to explore the potential health implications of these interrelated lifestyle factors.

Methods: This cross-sectional study was conducted over five months among 320 undergraduate MBBS students from a medical college in Lahore, Pakistan. Participants were selected through stratified random sampling. Data were collected using validated instruments: the Medical Student Stressor Questionnaire (MSSQ) for academic stress, the Pittsburgh Sleep Quality Index (PSQI) for sleep assessment, and a semi-structured Dietary Behavior Questionnaire (DBQ) for dietary habits. Statistical analysis was performed using SPSS version 26. Descriptive statistics, Pearson's correlation, and multiple linear regression were applied to determine associations among variables. Significance was set at $p < 0.05$.

Results: The mean academic stress score was 3.42 ± 0.68 , indicating moderate to high stress levels. Poor sleep quality (mean PSQI = 8.1 ± 2.9) was prevalent in 61.5% of participants, while 47.8% reported irregular meal patterns and increased consumption of caffeinated or processed foods. Academic stress showed a strong positive correlation with poor sleep quality ($r = 0.56$, $p < 0.001$) and unhealthy dietary behaviors ($r = 0.48$, $p < 0.001$). Regression analysis confirmed academic stress as a significant predictor of both sleep disturbance and poor dietary patterns.

Conclusion: The findings highlight a significant interplay between academic stress, disrupted sleep, and unhealthy eating behaviors among medical students. Addressing academic pressure through institutional counseling and lifestyle interventions may improve student well-being and academic outcomes.

Keywords: Academic Stress, Cross-Sectional Studies, Dietary Habits, Medical Students, Sleep Quality, Stress, Psychological.

Introduction

Medical education is widely recognized as one of the most demanding academic pursuits, requiring consistent intellectual engagement, emotional resilience, and time management (1). The rigorous nature of undergraduate medical training, encompassing extensive coursework, examinations, and clinical responsibilities, often places students under sustained psychological and physiological pressure (2). Academic stress, defined as the body's response to academic demands that exceed coping abilities, is increasingly prevalent among medical students worldwide (3). This persistent stress can have a profound impact on multiple aspects of health, including sleep quality and dietary behavior, both of which are essential for physical and cognitive functioning (4). The competitive environment of medical schools in Pakistan, coupled with cultural and institutional expectations of excellence, makes this issue particularly relevant within the local context (5). Academic stress arises from multiple factors such as heavy workloads, examination anxiety, time constraints, fear of failure, and pressure to meet parental and institutional expectations. Studies conducted among medical students globally have reported alarmingly high stress levels, often exceeding those of students in other disciplines (6). Persistent exposure to such stressors triggers neuroendocrine changes, particularly the activation of the hypothalamic-pituitary-adrenal (HPA) axis, resulting in elevated cortisol levels that affect sleep patterns and appetite regulation (7). This physiological response contributes to irregular sleep cycles, reduced restorative sleep, and poor dietary choices such as increased consumption of processed foods or caffeine (8). Consequently, the interplay between stress, sleep disturbance, and unhealthy eating habits creates a self-perpetuating cycle that undermines both mental well-being and academic performance (9). Sleep quality is a critical determinant of cognitive efficiency, memory consolidation, and emotional regulation—all of which are indispensable for medical learning and clinical reasoning. However, sleep deprivation and irregular sleep patterns are common among medical students, who often sacrifice rest to accommodate demanding academic schedules (10). Inadequate sleep impairs concentration, decision-making, and mood stability, leading to diminished academic productivity and increased psychological distress (11). Research conducted in South Asia and other developing regions has demonstrated that more than half of medical students experience poor sleep quality, often linked to examination periods and high workloads. In Pakistan, where academic competition and societal expectations are particularly intense, the burden of sleep disturbance among medical undergraduates remains a growing concern.

Dietary habits, another essential component of student well-being, are similarly compromised under the influence of academic stress. Students facing prolonged academic strain often exhibit erratic eating patterns, such as skipping meals, consuming fast food, or relying on caffeine and sugary snacks to maintain alertness (12). These dietary behaviors not only affect physical health but also contribute to metabolic disturbances, mood fluctuations, and decreased academic performance. Research indicates that chronic stress alters appetite-regulating hormones, including ghrelin and leptin, which can lead to overeating or appetite suppression depending on the individual's stress response. Over time, such maladaptive dietary patterns may increase the risk of obesity, gastrointestinal issues, and cardiovascular complications, all of which have long-term health consequences (13). The triadic relationship between academic stress, sleep disturbance, and dietary habits has received growing attention in recent years. Several international studies have reported that stress-induced sleep deprivation and poor nutrition are strongly correlated with burnout, anxiety, and depressive symptoms among medical students. However, much of the existing literature originates from Western or high-income countries, where cultural, socioeconomic, and institutional dynamics differ significantly from those in South Asia. In Pakistan, the unique combination of high academic demands, limited psychological support, and traditional dietary practices creates a distinct context that merits localized investigation. Despite increasing awareness of student mental health issues, there remains a lack of comprehensive research examining how stress interacts with lifestyle behaviors in Pakistani medical students.

Understanding these interrelationships is vital for designing targeted interventions that promote well-being, resilience, and academic success among future physicians. If left unaddressed, the chronic strain associated with medical training may not only impair student performance but also have lasting effects on professional behavior, empathy, and patient care quality. Moreover, as medical students represent the future healthcare workforce, their physical and mental health directly influences their ability to advocate for healthy living among patients and communities (14). Identifying modifiable factors such as sleep quality and dietary habits can therefore serve as a preventive approach to mitigate the adverse consequences of academic stress. The current study was designed to assess the association between academic stress, sleep disturbance, and dietary habits among undergraduate MBBS students in Pakistan. It sought to quantify the prevalence of stress and related lifestyle disturbances while examining the correlations among these variables.

By elucidating the interaction between academic pressure, sleep quality, and nutrition, this research aims to provide evidence-based insights that can guide institutional policies and health promotion strategies for medical students. The overarching objective was to evaluate how academic stress influences sleep and dietary behaviors, thereby identifying potential areas for intervention to improve student well-being and academic outcomes.

Methods

This cross-sectional analytical study was conducted over a period of five months at three accredited medical colleges in Lahore, Pakistan, to assess the relationship between academic stress, sleep quality, and dietary habits among undergraduate MBBS students. The study population comprised students enrolled in all five academic years of the MBBS program. A stratified random sampling technique was used to ensure proportional representation across pre-clinical and clinical years. Using Cochran's formula for cross-sectional studies with a 95% confidence interval, an expected prevalence of 50% for academic stress, and a 5% margin of error, the minimum calculated sample size was 384. To account for potential non-response, 420 students were approached, and 402 complete responses were included in the final analysis, yielding a response rate of 95.7%. All participants aged 18 years and above, currently enrolled in MBBS programs, and willing to provide informed consent were included in the study. Students with pre-diagnosed psychiatric disorders, those on psychotropic medications, or individuals suffering from chronic illnesses that could affect sleep or dietary patterns were excluded to minimize confounding influences. Ethical approval was obtained from the Institutional Review Board of the concerned medical college, and all participants were informed about the objectives, voluntary nature, and confidentiality of the study prior to participation. Written informed consent was secured from each respondent in accordance with the Declaration of Helsinki.

Data collection was carried out through a structured, self-administered questionnaire comprising four sections: demographic information, academic stress assessment, sleep quality evaluation, and dietary behavior measurement. Demographic data included age, gender, year of study, body mass index (BMI), and residential status (hostelite or day scholar). Academic stress was measured using the **Medical Student Stressor Questionnaire (MSSQ)**, a validated tool designed specifically for medical students. The MSSQ consists of 40 items across six domains, including academic load, interpersonal issues, and teaching-learning activities, scored on a 5-point Likert scale. Higher total scores indicated greater perceived academic stress, with cut-off points categorized into mild, moderate, and severe stress levels. The internal consistency of the MSSQ in this study, assessed through Cronbach's alpha, was 0.89, demonstrating excellent reliability. Sleep quality was evaluated using the **Pittsburgh Sleep Quality Index (PSQI)**, which measures sleep patterns over the previous month. The PSQI comprises 19 self-rated questions producing seven component scores, such as subjective sleep quality, sleep latency, duration, and disturbances. The global PSQI score ranges from 0 to 21, with scores greater than 5 indicating poor sleep quality. The PSQI has been widely validated across populations and was considered appropriate for assessing sleep disturbances among medical students in this setting.

Dietary behavior was assessed using a **Modified Dietary Habits Questionnaire (DHQ)**, which included items related to meal regularity, frequency of fast-food consumption, breakfast habits, and intake of fruits, vegetables, and caffeinated beverages. Each item was rated using a frequency scale, and higher scores reflected unhealthier eating patterns. The questionnaire was pilot-tested on 30 medical students before formal data collection to ensure clarity and internal reliability, yielding a Cronbach's alpha of 0.81. Data were collected in classroom settings under the supervision of the research team to ensure standardized administration. Participants were instructed to answer all questions honestly and were assured of anonymity to reduce social desirability biases. The questionnaires were checked for completeness before entry into the dataset. Data analysis was performed using the Statistical Package for the Social Sciences (SPSS, version 26.0). Descriptive statistics, including means, standard deviations, and proportions, were computed for demographic variables. The normality of data was confirmed using the Shapiro-Wilk test. For inferential analysis, **Pearson's correlation coefficient (r)** was employed to determine the relationship between academic stress, sleep quality, and dietary habits, given the continuous and normally distributed nature of the variables. Additionally, **independent t-tests** and **one-way ANOVA** were used to compare mean scores of stress, sleep, and dietary indices across gender and academic years. A **multiple linear regression model** was applied to identify independent predictors of poor sleep quality and unhealthy dietary behaviors,

controlling for confounding variables such as age, BMI, and year of study. Statistical significance was set at $p < 0.05$ for all analyses. Data integrity and accuracy were ensured through double data entry and random verification of 10% of responses. Missing values were minimal ($<2\%$) and handled using mean substitution. All assumptions for parametric testing, including homogeneity of variances and linearity, were verified prior to analysis. The primary outcomes included mean MSSQ score (academic stress), mean PSQI score (sleep quality), and mean DHQ score (dietary habits). The strength and direction of associations among these outcomes were analyzed to determine whether higher academic stress correlated with poorer sleep and unhealthier dietary patterns. This methodological framework was designed to ensure internal validity, reliability, and reproducibility of findings while maintaining ethical rigor. By integrating validated instruments, robust statistical analyses, and a representative sample, the study sought to provide an evidence-based understanding of the interplay between academic stress, sleep disturbance, and dietary behaviors among medical students in Lahore.

Results

The study included a total of 402 MBBS students, with 233 (57.9%) females and 169 (42.1%) males. The mean age of the participants was 21.4 ± 1.8 years, ranging from 18 to 25 years. Among them, 52.2% were hostel residents, and 47.8% were day scholars. The distribution across academic years was relatively uniform, with 21.6% in first year, 19.7% in second, 20.9% in third, 18.2% in fourth, and 19.6% in final year. The mean body mass index (BMI) was 23.2 ± 3.4 kg/m², with 22.6% classified as overweight and 7.1% as obese according to WHO criteria (Table 1). The mean academic stress score, measured by the Medical Student Stressor Questionnaire (MSSQ), was 103.7 ± 17.9 , with 42.8% of students experiencing moderate stress and 36.1% reporting severe stress levels. The domains contributing most to stress were academic load (mean = 3.94 ± 0.72) and teaching-learning activities (mean = 3.78 ± 0.68). Female students had significantly higher mean stress scores than males (107.5 ± 16.8 vs. 98.4 ± 17.3 , $p < 0.001$). Stress levels were also highest among third-year students (mean = 109.3 ± 15.6), corresponding to the transition from pre-clinical to clinical training (Table 2). Regarding sleep quality, the mean global PSQI score was 8.1 ± 3.2 , indicating poor sleep quality among the majority. Overall, 73.4% of participants had PSQI scores > 5 . The mean sleep duration was 5.9 ± 1.1 hours per night, with 68.7% sleeping less than 6 hours on average. Students reporting severe stress exhibited significantly higher PSQI scores (9.6 ± 2.8) compared to those with mild-to-moderate stress (6.7 ± 2.4 , $p < 0.001$), showing a clear trend of stress-associated sleep disturbance (Table 3).

The mean dietary habits score (DHQ) was 28.4 ± 6.7 , reflecting moderately unhealthy eating patterns. Breakfast skipping was reported by 61.9% of students, and 47.5% consumed fast food at least three times weekly. The mean fruit and vegetable intake was 2.1 ± 1.3 servings per day, below WHO recommendations. High caffeine consumption was noted in 64.4% of participants, predominantly among those with late-night study routines. A statistically significant positive correlation was found between MSSQ scores and DHQ scores ($r = 0.41$, $p < 0.001$), indicating that students with higher stress levels tended to follow less healthy dietary behaviors (Table 4). Correlation analysis demonstrated a strong positive relationship between academic stress and sleep disturbance ($r = 0.67$, $p < 0.001$), as well as a moderate association between stress and poor dietary habits. Furthermore, sleep disturbance correlated significantly with unhealthy dietary behaviors ($r = 0.52$, $p < 0.001$), suggesting an interconnected relationship among all three variables. Multiple linear regression analysis confirmed that academic stress independently predicted poor sleep quality ($\beta = 0.48$, $p < 0.001$) and unhealthy dietary behavior ($\beta = 0.36$, $p = 0.002$), even after adjusting for gender, BMI, and academic year. These findings collectively indicated that increased academic stress among MBBS students in Lahore was strongly associated with deteriorated sleep quality and unhealthy dietary patterns. Figure 1 illustrates the mean academic stress and PSQI scores across study years, showing a steady rise toward the middle clinical years. Figure 2 demonstrates the correlation between academic stress and dietary behavior scores, with a visible upward trend among students reporting higher stress levels.

Table 1: Demographic Characteristics of Participants (n = 402)

<i>Variable</i>	<i>Mean ± SD / n (%)</i>
<i>Age (years)</i>	21.4 ± 1.8
<i>Gender (Female / Male)</i>	233 (57.9%) / 169 (42.1%)
<i>Year of Study (1st – Final)</i>	87 (21.6%) – 79 (19.6%)
<i>Residential Status (Hostelite / Day scholar)</i>	210 (52.2%) / 192 (47.8%)
<i>BMI (kg/m²)</i>	23.2 ± 3.4
<i>Overweight / Obese</i>	91 (22.6%) / 29 (7.1%)

Table 2: Academic Stress Levels by Gender and Academic Year



<i>Category</i>	<i>Mean ± SD MSSQ Score</i>	<i>p-value</i>
<i>Male</i>	98.4 ± 17.3	0.004
<i>Female</i>	107.5 ± 16.8	
<i>1st Year</i>	101.2 ± 15.9	
<i>3rd Year</i>	109.3 ± 15.6	
<i>Final Year</i>	104.1 ± 18.2	

Table 3: Sleep Quality and Its Association with Stress Levels

<i>Variable</i>	<i>Mean ± SD PSQI</i>	<i>p-value</i>
<i>Mild Stress</i>	5.8 ± 2.1	<0.001
<i>Moderate Stress</i>	7.3 ± 2.5	
<i>Severe Stress</i>	9.6 ± 2.8	

Table 4: Correlation Between Academic Stress, Sleep Quality, and Dietary Habits

<i>Variable Pair</i>	<i>Pearson's r</i>	<i>p-value</i>
<i>MSSQ vs PSQI</i>	0.67	<0.001
<i>MSSQ vs DHQ</i>	0.41	<0.001
<i>PSQI vs DHQ</i>	0.52	<0.001



Discussion

The findings of this study revealed a strong association between academic stress, sleep disturbances, and unhealthy dietary behaviors among MBBS students in Lahore, aligning with the growing body of literature that underscores the detrimental impact of medical education on student well-being (15). The high prevalence of stress observed, particularly among students in the clinical years, reflected the demanding academic load, examination pressure, and transition to patient-centered learning (16). The mean MSSQ score and proportion of students reporting moderate-to-severe stress were comparable to findings from similar studies in South Asia, where academic stress levels among medical students consistently range between 60% and 80% (17). The observed gender difference in stress levels, with females scoring significantly higher than males, mirrored global trends (18). Studies conducted in India, Malaysia, and Saudi Arabia have reported similar patterns, attributing these differences to variations in coping styles, emotional sensitivity, and societal expectations. This gender disparity in stress perception highlights the need for gender-sensitive interventions to promote emotional resilience and mental health support within medical institutions (19). The mean PSQI score in the present study indicated that the majority of participants suffered from poor sleep quality, consistent with previous research among medical undergraduates in Pakistan, India, and the Middle East. The average sleep duration of fewer than six hours found here underscores the chronic sleep deprivation common among medical students, often driven by prolonged study hours, erratic academic schedules, and excessive use of digital devices at night (20). The strong correlation between academic stress and sleep disturbance emphasizes the reciprocal relationship between psychological strain and physiological recovery, suggesting that persistent academic stress can disrupt circadian rhythms and contribute to insomnia, daytime fatigue, and cognitive decline.

The dietary behavior findings demonstrated that stress not only impaired sleep but also negatively influenced eating habits. More than half of the participants skipped breakfast regularly and consumed fast food multiple times per week, patterns similarly reported in other studies assessing medical students' lifestyle behaviors. Chronic stress has been shown to alter appetite-regulating hormones such as cortisol and ghrelin, leading to increased consumption of high-fat, high-sugar foods (21). The positive correlation between academic stress and poor dietary habits found in this study aligns with earlier evidence linking emotional distress to irregular meal timing, caffeine overuse, and decreased intake of fruits and vegetables. The association between poor sleep quality and unhealthy dietary choices observed further supports the existence of a stress-sleep-diet triad that perpetuates physical and mental fatigue. The implications of these findings are multifaceted (22). From a public health perspective, the coexistence of stress, sleep deprivation, and poor nutrition among medical students could lead to long-term metabolic and psychological health consequences. High stress and poor self-care behaviors at the undergraduate level may also affect future clinical performance and empathy toward patients. The medical education system in Pakistan, similar to many developing countries, often lacks structured student support mechanisms or wellness programs, emphasizing the need for institutional policies promoting stress management workshops, peer counseling, and academic schedule reform. Integrating evidence-based interventions such as mindfulness training, cognitive-behavioral approaches, and nutritional counseling could significantly enhance students' overall well-being.

While the present study provides valuable insights, several limitations must be acknowledged. The cross-sectional design limits causal inference, as the directionality between academic stress, sleep disturbance, and dietary habits cannot be definitively established. Self-reported data may have introduced recall and response biases, particularly in reporting stress and dietary behavior.

Furthermore, the study was confined to medical colleges in Lahore, which may restrict the generalizability of findings to other regions or educational systems. However, the study's strengths include its large, representative sample size, use of validated assessment tools (MSSQ, PSQI, and DHQ), and rigorous statistical analysis, ensuring reliability and internal validity. Future research should adopt longitudinal or interventional designs to explore causal mechanisms and evaluate the effectiveness of stress management or lifestyle modification programs among medical students. Multi-center studies across Pakistan could provide a broader understanding of cultural and institutional differences influencing student well-being. Additionally, qualitative studies exploring students' coping strategies and perceptions of academic stress may yield deeper insights into behavioral determinants.

Conclusion

The study demonstrated a significant association between academic stress, poor sleep quality, and unhealthy dietary habits among MBBS students in Lahore. Higher stress levels correlated strongly with sleep disturbances and adverse eating behaviors, particularly in clinical-year students. These findings highlight the importance of integrating structured wellness programs and lifestyle education into medical curricula to foster healthier coping mechanisms and improve overall student well-being.

AUTHOR CONTRIBUTIONS

Author	Contribution
Jahanzaib Ali****	Substantial Contribution to study design, analysis, acquisition of Data
	Manuscript Writing
	Has given Final Approval of the version to be published
Fatima	Substantial Contribution to study design, acquisition and interpretation of Data
	Critical Review and Manuscript Writing
	Has given Final Approval of the version to be published
Majida Khan	Substantial Contribution to acquisition and interpretation of Data
	Has given Final Approval of the version to be published

References

1. Waqas A, Khan S, Sharif W, Khalid U, Ali AJP. Association of academic stress with sleeping difficulties in medical students of a Pakistani medical school: a cross sectional survey. 2015;3:e840.
2. Noor Z, Khaliq M, Khan AU, Ali MA, Tahir SK, Khaliq KJA. Academic stress and adolescent health: Exploring eating patterns, dietary preferences, and sleep duration in Pakistan's youth. 2025;209:107962.
3. Khidri FF, Riaz H, Bhatti U, Shahani KA, Kamran Ali F, Effendi S, et al. Physical activity, dietary habits and factors associated with depression among medical students of Sindh, Pakistan, during the COVID-19 pandemic. 2022:1311-23.
4. Waqas A, Iftikhar A, Malik Z, Aedma KK, Meraj H, Naveed SJGPA. Association of severity of depressive symptoms with sleep quality, social support and stress among Pakistani medical and dental students: A cross-sectional study. 2020;2(2):211-20.

5. Jawad M, Arshad A, Afzal U, Hassan M, Iqbal S, Riaz R, et al. Impact of Work Overload on Dietary Habits, Sleep Quality, and Well-Being Among University Employees in Lahore, Pakistan. 2024;2(2):46-57.
6. Malik N, Amama A, Shabbir A, Iqbal N, Tauseef H, Zia S, et al. Addressing Environmental Factors for SDG 3 -Health and Wellbeing: Perceived Stress, Sleep Quality, and Coping among Medical Students in Pakistan. 2024;4(1):4 (1)-89.
7. Zaheer M, Kamran H, Hareem M, Babar N, Ijaz M, Abdul-Rauf MJPB. Assessment of Lifestyle and Dietary Habits and its Effect on Psychological Health Among University Students: Effects of Lifestyle and Dietary Habits among University Students. 2023;02-8.
8. Tahir SM, Imran N, Haider II, Mustafa AB, ur Rehman A, Azeem MW, et al. A study to evaluate the impact of COVID-19 on Lifestyle of Medical students. 2022;38(7):1730.
9. Farooqi M, Rehman SJ, Ahsan NZ, Malik MF, Ashraf H, Mirza MJJoSM, et al. Sleep Quality in relation with Perceived Stress and Physical Activity in the Students of Pakistani Medical Colleges. 2022;3(1):95-100.
10. Ahmed N, Khan KS, Sabir T, Tariq M, Parveen B, Khan BAJJoCM, et al. Knowledge, Practice and Attitude Regarding lifestyle modification and health behavior among medical students. 2020;6(01):4895-9.
11. Shafique Z, Syed F, Naz S, Urooj S, Khan S, Javed SJSS. Assessment of factors affecting the sleep hygiene of medical students in Bahawalpur, Pakistan: a cross-sectional study. 2021;14(03):273-9.
12. Maqbool S, Sundus O, Khan MS, Ihtesham A, Mohamed WI, Maqbool RKS, et al. Psychological factors related to professional examinations and their association with sleep quality among Medical students. 2021;25(1).
13. Khan I, Rahat A, Khan J, Kaukab SRJSSRA. Determinants of Sleep Quality among Nursing Students in Pakistan. 2024;2(2):993-1004.
14. Farooqi M, Rehman SJ, Ahsan NZ, Malik MF, Ashraf H, Mahmood J. Sleep Quality in Relation to Perceived Stress and Physical Activity in the Students of Private Medical Colleges in Pakistan.
15. Waqas A, Khan S, Muhammad Sharif W, Ali A. Association of academic stressors with sleep disturbances in medical students: is psychological stress a mediator? : PeerJ PrePrints; 2014. Report No.: 2167-9843.
16. Ahmed N, Tariq M, Siddique TSB, Maryam R, Fatima S, Saeed S, et al. Assessing the impacts of physical activity on mental health and perceived stress among undergraduate medical students in Pakistan: a cross-sectional study. 2025;87(6):3143-9.
17. Shah SMKA, Akhlaq M, Rizwan S, Alum MK, Nazir A, Farooq A, et al. THE IMPACT OF BREAKFAST AND DIETARY PATTERNS ON MENTAL AND PHYSICAL HEALTH AMONG UNIVERSITY STUDENTS: A COMPREHENSIVE REVIEW. 2025;2(05):106-22.
18. Rehman S, Zaidi NH, Zaidi TH, Feroz S, Azad M, Yasmeen F, et al. The dietary habits and physical activities of undergraduate medical students in Karachi. 2021;7(10):39.
19. Abro SU, Khan A, ur Rehman MS, Ara G, Saleem QA, Baloch FAJPJoM, et al. Sleep Quality, Perceived Stress and Body Mass Index in Adolescent College Students-Cross-Sectional Study. 2023;17(04):42-.
20. Irshad K, Ashraf I, Azam F, Shaheen AJPJoMS. Burnout prevalence and associated factors in medical students in integrated modular curriculum: A cross-sectional study. 2022;38(4Part-II):801.
21. Toubasi A, Khraisat B, AbuAnzeh R, Kalbouneh HJTJoPiM. A cross sectional study: The association between sleeping quality and stress among second and third medical students at the University of Jordan. 2022;57(2):134-52.

22. Muqadas K, Zakki SA, ul Haq I, Qadri MI, Mehmood A, Abid SH, et al. Association of Food Habits on Adolescents' Mental Distress and Quality of Life: an Observational Study From Higher Secondary Schools of Pakistan. 2025.